



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2020

---

## Structure and unique mechanical aspects of nuclear lamin filaments

Tenga, Rafael ; Medalia, Ohad

**Abstract:** The major constituent of the nuclear envelope is the nuclear lamina. A fibrous meshwork of lamin filaments spanning underneath the nuclear membrane provides mechanical support to the nucleus. Lamins, type V intermediate filament proteins, are also involved in many other nuclear activities such as DNA repair and transcription. Recent structural studies provide new insight into how lamins assemble into 3.5nm thick filaments, which is in contrast with cytoplasmic intermediate filaments. The thinnest component of the cell cytoskeleton exhibits surprising mechanical properties. Here, we critically review and discuss structural and mechanical aspects of single lamin filaments.

DOI: <https://doi.org/10.1016/j.sbi.2020.06.017>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-192748>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Tenga, Rafael; Medalia, Ohad (2020). Structure and unique mechanical aspects of nuclear lamin filaments. *Current Opinion in Structural Biology*, 64:152-159.

DOI: <https://doi.org/10.1016/j.sbi.2020.06.017>



# Structure and unique mechanical aspects of nuclear lamin filaments

Rafael Tenga and Ohad Medalia

The major constituent of the nuclear envelope is the nuclear lamina. A fibrous meshwork of lamin filaments spanning underneath the nuclear membrane provides mechanical support to the nucleus. Lamins, type V intermediate filament proteins, are also involved in many other nuclear activities such as DNA repair and transcription. Recent structural studies provide new insight into how lamins assemble into ~3.5 nm thick filaments, which is in contrast with cytoplasmic intermediate filaments. The thinnest component of the cell cytoskeleton exhibits surprising mechanical properties. Here, we critically review and discuss structural and mechanical aspects of single lamin filaments.

## Address

Department of Biochemistry, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

Corresponding author: Medalia, Ohad ([omedalia@bioc.uzh.ch](mailto:omedalia@bioc.uzh.ch))

**Current Opinion in Structural Biology** 2020, **64**:152–159

This review comes from a themed issue on **Cryo electron microscopy**

Edited by **Sara Sandin** and **Daniela Rhodes**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 15th August 2020

<https://doi.org/10.1016/j.sbi.2020.06.017>

0959-440X/© 2020 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

The inner and outer nuclear membranes (INM and ONM, respectively) provide a physical barrier that is perforated by the nuclear pore complex (NPC), an enormous macromolecular complex that provide the sole gateways in and out of the nucleus [1,2]. Attached to the INM and interacting with NPCs, the nuclear lamina [3–6] provides a layer of filamentous scaffolding that is anchored to the membrane as well as to the NPCs. Composed of lamins and lamin-associated proteins, which include emerin and LBR among others [7,8], the 10–30 nm thick nuclear lamina is found ubiquitously in differentiated cells.


Over the years it has been shown that lamins play a role in many nuclear functions, for example, transcription regulation, replication, DNA repair and chromatin organization [7,9–12]. Four main lamin isoforms are found in mammals. In humans, they are encoded by

the *LMNA*, *LMNB1*, *LMNB2* genes to express lamin A/C, lamin B1 and lamin B2, respectively. While the B-type lamins are expressed in all types of mammalian cells, the expression of A-type lamins is developmentally regulated and appears only during cell differentiation [13,14]. Notably, the expression of A-type and B-type lamins is highly variable and depends on the cell type and differentiation stage [15]. Surprisingly, recent studies showed that differentiated cells lacking B-type lamins are viable [16]. In differentiated cells, both types of lamins are localized to the nuclear periphery; however, small amounts of A-type lamins are also found in the nucleoplasm and are suggested to have a major functional impact on chromatin organization and gene regulation [17].

Mutations in lamin A/C have been directly associated with >15 different diseases, termed laminopathies [18]. Laminopathies are highly varied and include muscular dystrophies [19], lipodystrophies [20] and progeria syndromes [21] to name a few. However, it is still not clear, how the single point mutations can lead to such a wide variety of diseases and it remains an open and heavily studied question [5]. Many of these mutations affect tissues under high mechanical load like muscles and heart [22,23]. This may suggest that mutations also alter the mechanical properties of lamins in addition to alteration of their structure and interaction with other proteins [24,25].

Lamins are classified as type V intermediate filament (IF) proteins and are considered to be the ancestors of cytoplasmic IFs [26]. Similar to other IF proteins, lamins contain a long rod domain which comprises four coiled-coil  $\alpha$ -helical segments, termed 1A, 1B, 2A and 2B, separated by flexible linkers (Figure 1a). This domain is flanked by a non-helical N-terminal head and C-terminal tail domain. The C-terminal domain hosts a nuclear localization signal (NLS), an immunoglobulin (Ig) like fold and a CaaX motif box (C, cysteine; a, aliphatic residue; X, any residue). Lamins are post-translationally modified at the CaaX motif by farnesylation of the cysteine residue (except for lamin C, which is devoid of the CaaX motif box). Lamin A is further cleaved before the -aaX sequence by a metallo-peptidase (Zmpste24 or endoprotease RAS converting enzyme 1 (RCE1)). Thus, only B-type lamins remain farnesylated, and therefore their interactions with the INM are retained. Furthermore, post-translational modifications (PTMs) of mature lamins include multiple

Figure 1



Nuclear lamins: proteins and assemblies. **(a)** The lamin protein-domains organization is conserved among species. The domains of lamin A/C, B1, B2 as well as the single *C. elegans* lamin the and frog oocyte LIII are shown. **(b)** Bacterially expressed *C. elegans* lamin assembled under low ionic strength buffers into ~8 nm thick filaments, imaged by cryo-electron tomography [46]. **(c)** Human lamin A assembled *in vitro* into paracrystalline arrays, imaged by cryo-electron tomography. These paracrystalline structures are not ordered enough to allow the structural determination of the lamin assemble. **(d)** A tomographic slice through the nuclear lamina of a nuclease treated mammalian nucleus reveals the lamin filaments meshwork. **(e)** A tomographic slice through a native nuclear lamina as observed by cryo-ET of cryo-FIB milling of a MEF nucleus, lamins filaments (arrow heads) and chromatin (arrow) are seen. **(f)** 2D averaging revealed structural classes of lamin filaments. A 3.5 nm thick rod domain (blue arrow head) decorated with globular Ig-folds (red arrow head) with a 20 nm repeat (adapted from [58]).

phosphorylations [27] that can promote lamin solubility and delocalization of lamin A into the nucleoplasm [28,29].

Recent studies showed that lamins are involved in mechano-signaling and have a major role in force transduction [30–32]. We hypothesize that these unique properties must be reflected in the structure of lamin proteins and subsequent filament assembly. In this review, we provide a summary of lamin structures and the mechanical properties of these intermediate filament proteins.

### Towards structural determination of lamins

Intermediate filaments assemble into dimers in solution. Lamin dimer analysis revealed the formation of a

parallel coiled-coil structure between two monomers using metal shadowing transmission electron microscopy (TEM) [33]. Lamin dimers further assemble by head-to-tail association into long polymers, which further associate laterally into the mature filaments [34]. These assembly steps were only identified at low resolution and are still awaiting to be resolved in high resolution.

Recent biochemical analysis of *in vitro* assembled lamin A dimers and tetramers suggested that the structure of lamin dimers is affected by the chemical environment during the assembly, for example, electrolyte compositions [35]. Linker regions within the lamin proteins can be compressed, leading to shorter repeat units. This implies that flexible linker regions allow sliding of

dimeric coiled-coil domains within tetrameric higher-order assemblies.

High-resolution structural determination is required to detect sliding of lamin dimers and tetramers. However, due to the length of lamin dimers, the coiled coil domain is ~50 nm while the tail and the head domains are not contributing to the filament scaffold, structural determination of lamin dimers is a challenging task [36]. Their tendency to rapidly polymerize makes full-length lamins as well as the intact coiled-coil domains difficult to approach in structural studies by x-ray crystallography [37]. A detailed atomic model of lamin proteins is still unavailable, however, a 'divide-and-conquer' crystallization strategy has provided insights into the organization of the lamins, by determining the crystal structure of (mostly) lamin A fragments [37,38]. The atomic structure of the Ig-fold of human lamin A/C was determined, exhibiting a globular two  $\beta$ -sheets structure [39,40]. The analysis of coiled-coil fragments provided insights into the intra-organization of lamin dimers [41].

The structures of lamin coiled-coil  $\alpha$ -helical segments containing 1A, 1B and small part of 2A exemplified the interactions between lamin coil 1B to form a tetrameric assembly were recently resolved by x-ray crystallography [42<sup>\*\*</sup>,43<sup>\*\*</sup>]. These studies revealed the mode of interactions between the 1B coiled-coil domain of dimers to form tetramers. These interactions suggested to be conserved in other IF proteins, for example, vimentin and keratins [43<sup>\*\*</sup>]. Surprisingly, it showed that the linkers positioned within the coiled-coil domain are folded into  $\alpha$ -helical structures, and form a continuum of the coiled-coil structure. Furthermore, it was suggested that interdimer contacts between helix 1 promotes the antiparallel interaction of helix 2 that may promote the longitudinal assembly of lamins [42<sup>\*\*</sup>,43<sup>\*\*</sup>]. However, high-resolution structural analysis of lamin assemblies is required in order to confirm these interactions in higher-order lamin assemblies.

Lamin assembly was studied *in vitro* and showed to be a hierarchical process. The chemical condition to assemble the *Caenorhabditis elegans* lamin into stable filaments are established, in contrast to recombinantly purified lamin from other organisms. Hence, these filaments can be analyzed by electron microscopy approaches (Figure 1b) [44–46]. A major disadvantage of the *in vitro* structural analysis of lamins is the tendency of the mammalian lamins to assembled into paracrystalline arrays (Figure 1c), which are probably not relevant in a biological context [34]. This observation suggests that assembly of lamin filaments requires-specific conditions or maybe some factors that are found at the nuclear envelope (NE) to provide the precise environment for homogeneous lamin filament assembly (Figure 1d–f).

## Architecture of lamins at the nuclear lamina

Cryo-electron tomography (cryo-ET) is a powerful technique to acquire structural insight into single non-repeating structures [47–49] and therefore became pivotal in cell biology [34,50,51]. This method enables structure determination of macromolecular complexes within a cell [52–55]. Recent technological developments allow to acquire a high-resolution snapshot of molecular processes at a specific point of time [56,57]. Therefore, cryo-ET is the method of choice to decipher the structure of lamin assemblies at close-to-native state.


Recently, we showed that lamin filaments assemble into 3.5 nm thick filaments (Figure 1f) within a ~14 nm thick meshwork layer beneath the nuclear membrane [58<sup>\*</sup>]. The filaments are both highly variable in length and exhibit a short persistence length of <200 nm which hints at a large flexibility. This persistence length means that lamin filaments are more bendable than any other components of the cytoskeleton, including other IF proteins. This physical characteristics of lamins represent their unique mechanical properties that are observed when lamin filaments are subjected to external forces [59,60<sup>\*</sup>].

In the nuclear lamina, the Ig-fold of lamins were seen as globular domains decorating the filaments every 20 nm. The different lamin isoforms could be partially identified using immunogold-labeling. It showed that A-type and B-type lamins form two separate meshworks, confirming previous observations obtained by structural illumination microscopy (3D-SIM) that lamin A, C, B1 and B2 form individual meshworks within the nuclear lamina [61]. An additional study using stochastic optical reconstruction microscopy (STORM) showed that the lamin meshworks are also spatially distinguished, with the lamin B1 meshwork lying closer to the membrane and the highest concentration of lamin A/C is found further towards the nucleoplasm [62<sup>\*</sup>].

The nuclear lamina contains an additional set of proteins and has been shown to tightly bind chromatin [63,64]. Therefore, the nuclear lamina plays a major role in the nuclear architecture and gene expression [63,65]. The heterochromatin domains, which bind to the lamina are referred to as lamina-associated domains (LADs) and are kept in a transcriptionally repressed state to maintain genome stability [66]. It was recently shown that the interaction of lamin B1 with chromatin is synchronized with the circadian clock [67]. The interactions between heterochromatin and lamin can be observed by means of cryo-ET (Figure 2). This image reveals that the chromatin is intertwined between nuclear lamins to form a direct interaction which can resist nuclease treatments. However, studying more native and intact nuclear samples is needed to decipher the complicated interactions between chromatin and lamins.



Figure 2



Visualizing lamin-chromatin interactions. (a) Applying nuclease treatment to lysed cells, as described in [58], retains heterochromatin and lamins, as revealed by visualizing HP1α and LaminA/C. (b) and (c) Segmented cryo-tomogram shows tight interactions between lamin filaments (yellow) and heterochromatin (blue).

The use of cryo-FIB (focused ion-beam) milling as a sample preparation procedure for cryo-ET allows to study any cellular structures inside the native environment of vitrified cells and multicellular samples [68,69]. By applying recent advances in cryo-FIB milling, in combination with cryo-ET [49,70,71<sup>••</sup>], it is possible to study structures in the nucleus [72–74]. FIB milling is used to create thin lamellas of cells or other structures which are too thick (>1 μm) and therefore unsuitable for native cryo-ET. Even subtomogram averaging approaches of nuclear structures like the NPC and nucleosome are feasible [73,75]. Therefore, not only lamins but all lamin-associated proteins, for example, the NPCs, the chromatin and the nuclear membranes can be studied. Thereby, combining cryo-FIB and cryo-ET allows us to study the structural interactions of the nuclear lamina with these cellular structures at high resolution. A typical view into the nuclear lamina as revealed by applying FIB-milling in conjunction with cryo-ET is shown in Figure 1e.

### Mechanical properties of nuclear lamin filaments


Lamins contribute to the mechanical stability of the nucleus, transduce mechanical forces and biochemical signals across the NE and regulate the organization of chromatin [30,76]. Studies performed on the nucleus and entire organisms are very informative but influenced by nuclear membranes, chromatin, and

surrounding cells, respectively. Such studies have assessed the changes to differing levels of lamins and NE proteins, and the underlying alterations of the physical properties [24,77]. Although chromatin plays a major role in nuclear mechanics [78], the question if lamins have special mechanical properties, is of major importance [79<sup>••</sup>].

A first glimpse into the mechanical properties of the lamins came from *in vitro* assembled macrofilaments that were subjected to external mechanical stress [59]. This macroscopic paracrystalline assemblies showed that lamins can withstand strong external forces. However, native A-type and B-type lamin filaments can only be assembled *in situ*. Such a study was recently conducted using a *Xenopus laevis* oocyte. The *X. laevis* oocyte lamin LIII revolutionized our understanding on the nuclear lamina since it was first visualized [80]. In this setup, a manually opened NE is visualized both by cryo-ET as well as by atomic force microscopy (AFM). Therefore, the mechanical properties of the lamin LIII can be directly measured within an *in situ* assembled lamina. Furthermore, the spread NE is free of chromatin, therefore only mechanical properties from the nuclear lamina are measured.

AFM-based force-extension (FE) measurements revealed that lamin filaments deform in a reversible manner under low loads (<500 pN) (Figure 3). However,

Figure 3



The lamin filaments under stress. A Model depicting the deformation of a lamin filament under external stress. Under low force the filaments are deformed by unfolding of the helical domains. This adaption to the force is reversible. When higher force is applied, the filaments stiffens by transitioning the  $\alpha$ -helical structures to  $\beta$ -sheets. This is an irreversible process that may lead to the filament breakage.

when force is increased ( $>2$  nN), lamins are stiffening substantially until the filament is failing. This suggests that lamins are a special material that is very elastic when a low force is induced but becomes a plastic stiff material when stronger forces are applied. Thus, lamins can react to force and change their behaviour dependant on the amount of force. Furthermore, molecular dynamics analysis suggested that the low force rearrangements are caused by reversible reorganization of the coiled-coil helical domains that may involve local unfolding. However, upon stronger forces lamins undergo a major irreversible transformation from an  $\alpha$ -helix to a  $\beta$ -sheet that make these filaments a stiff material. This suggests that upon strong external forces lamins are 'sacrificed' by irreversible unfolding to physically protect the nucleus (Figure 3). Mechanical protection of the nucleus by the lamin meshwork are integrated at each level of the hierarchical construction of lamins, from the basic building block of the coiled-coil domains up to the higher order meshwork. Lamin filaments are extensible, strong and tough, similar to natural silk and superior to the synthetic polymer Kevlar®. Although the study was conducted on the LIII lamins, structural analysis by cryo-ET showed the similarity to mammalian lamins and the nuclear lamina meshwork of fibroblasts (Figure 3).

## Concluding remarks

Lamin filaments are essential components of the cell nucleus. These thin IF proteins,  $\sim 3.5$  nm in diameter, are assembled underneath the nuclear envelope although a pool of A-type lamins is located in the nucleoplasm and was found to interact with specific binding partners, such as LAP2 $\alpha$  [17]. These lamin A/C structures are more dynamic than the ones at the peripheral lamina and likely have unique structural assembly states and functions, fundamentally different from those of the peripheral lamina. However, the *in vitro* assembly conditions for mammalian lamins have not been found yet. Surprisingly, lamin filaments have unique physical properties that may have evolved to fulfill their major task — to provide unique mechanical properties to the nucleus. High-resolution information is crucial to unveil the alteration of the nuclear lamins organization at the lamins, nuclear interior and in health and disease, but also when nuclei experience external forces, for example, when cell pass through narrow passages [24,25]. Changes in the lamina during the cell cycle may also require the correlation between structural analysis and cell stages. With recent advancements in cryo-ET as well as complementary modalities, these fundamental aims are within reach.

## Conflict of interest statement

Nothing declared.

## Funding

Funding was received for this work.

## Acknowledgements

We thank Rebecca de Leeuw for critical reading of the manuscript. This work was funded by grants from the Swiss National Science Foundation Grant (SNSF 31003A\_179418) and the Mäxi Foundation. We thank the Center for Microscopy and Image Analysis at the University of Zurich (ZMB).

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as

- of special interest
- of outstanding interest

1. Grossman E, Medalia O, Zwerger M: **Functional architecture of the nuclear pore complex**. *Annu Rev Biophys* 2012, **41**:557-584.
2. Hampoelz B, Andres-Pons A, Kastiris P, Beck M: **Structure and assembly of the nuclear pore complex**. *Annu Rev Biophys* 2019, **48**:515-536.
3. Burke B, Stewart CL: **The nuclear lamins: flexibility in function**. *Nat Rev Mol Cell Biol* 2013, **14**:13-24.
4. Dechat T, Adam SA, Taimen P, Shimi T, Goldman RD: **Nuclear lamins**. *Cold Spring Harb Perspect Biol* 2010, **2**:a000547.
5. Tatti M, Medalia O: **Insight into the functional organization of nuclear lamins in health and disease**. *Curr Opin Cell Biol* 2018, **54**:72-79.

6. Kittisopikul M, Shimi T, Tatli M, Tran JR, Zheng Y, Medalia O, Jaqaman K, Adam SA, Goldman RD: **Nuclear lamins A/C and B1 provide a structural framework that organizes and anchors nuclear pore complexes.** *bioRxiv* 2020, **2020** 2004.2003.022798.
  7. de Leeuw R, Gruenbaum Y, Medalia O: **Nuclear lamins: thin filaments with major functions.** *Trends Cell Biol* 2018, **28**:34-45.
  8. Cutler JA, Wong X, Hoskins VE, Gordon M, Madugundu AK, Pandey A, Reddy KL: **Mapping the micro-proteome of the nuclear lamina and lamin associated domains.** *bioRxiv* 2019. 828210.
  9. Mattout A, Cabcianca DS, Gasser SM: **Chromatin states and nuclear organization in development—a view from the nuclear lamina.** *Genome Biol* 2015, **16**:174.
  10. Kennedy BK, Barbie DA, Classon M, Dyson N, Harlow E: **Nuclear organization of DNA replication in primary mammalian cells.** *Genes Dev* 2000, **14**:2855-2868.
  11. Markiewicz E, Tilgner K, Barker N, van de Wetering M, Clevers H, Dorobek M, Hausmanowa-Petrusewicz I, Ramaekers FC, Broers JL, Blankestijn WM *et al.*: **The inner nuclear membrane protein emerin regulates beta-catenin activity by restricting its accumulation in the nucleus.** *EMBO J* 2006, **25**:3275-3285.
  12. Maynard S, Keijzers G, Akbari M, Ezra MB, Hall A, Morevati M, Scheibye-Knudsen M, Gonzalo S, Bartek J, Bohr VA: **Lamin A/C promotes DNA base excision repair.** *Nucleic Acids Res* 2019, **47**:11709-11728.
  13. Kim Y, Sharov AA, McDole K, Cheng M, Hao H, Fan CM, Gaiano N, Ko MS, Zheng Y: **Mouse B-type lamins are required for proper organogenesis but not by embryonic stem cells.** *Science* 2011, **334**:1706-1710.
  14. Constantinescu D, Gray HL, Sammak PJ, Schatten GP, Csoka AB: **Lamin A/C expression is a marker of mouse and human embryonic stem cell differentiation.** *Stem Cells* 2006, **24**:177-185.
  15. Shin JW, Spinler KR, Swift J, Chasis JA, Mohandas N, Discher DE: **Lamins regulate cell trafficking and lineage maturation of adult human hematopoietic cells.** *Proc Natl Acad Sci U S A* 2013, **110**:18892-18897.
  16. Kim Y, Zheng X, Zheng Y: **Proliferation and differentiation of mouse embryonic stem cells lacking all lamins.** *Cell Res* 2013, **23**:1420-1423.
  17. Naeter N, Ferraioli S, Foisner R: **Lamins in the nuclear interior - life outside the lamina.** *J Cell Sci* 2017, **130**:2087-2096.
  18. Wong X, Stewart CL: **The laminopathies and the insights they provide into the structural and functional organization of the nucleus.** *Annu Rev Genomics Hum Genet* 2020.
  19. Bonne G, Di Barletta MR, Varnous S, Becane HM, Hammouda EH, Merlini L, Muntion F, Greenberg CR, Gary F, Urtizberea JA *et al.*: **Mutations in the gene encoding lamin A/C cause autosomal dominant Emery-Dreifuss muscular dystrophy.** *Nat Genet* 1999, **21**:285-288.
  20. Caron M, Auclair M, Donadille B, Bereziat V, Guerci B, Laville M, Narbonne H, Bodemer C, Lascols O, Capeau J *et al.*: **Human lipodystrophies linked to mutations in A-type lamins and to HIV protease inhibitor therapy are both associated with prelamin A accumulation, oxidative stress and premature cellular senescence.** *Cell Death Diff* 2007, **14**:1759-1767.
  21. De Sandre-Giovannoli A, Bernard R, Cau P, Navarro C, Amiel J, Boccaccio I, Lyonnet S, Stewart CL, Munnich A, Le Merrer M *et al.*: **Lamin A truncation in Hutchinson-Gilford progeria.** *Science* 2003, **300**:2055.
  22. Earle AJ, Kirby TJ, Fedorchak GR, Isermann P, Patel J, Iruvanti S, Moore SA, Bonne G, Wallrath LL, Lammerding J: **Mutant lamins cause nuclear envelope rupture and DNA damage in skeletal muscle cells.** *Nat Mater* 2020, **19**:464-473.
  23. Hah J, Kim DH: **Deciphering nuclear mechanobiology in laminopathy.** *Cells* 2019, **8**.
  24. Harada T, Swift J, Irianto J, Shin JW, Spinler KR, Athirasala A, Diegmiller R, Dingal PC, Ivanovska IL, Discher DE: **Nuclear lamin stiffness is a barrier to 3D migration, but softness can limit survival.** *J Cell Biol* 2014, **204**:669-682.
  25. McGregor AL, Hsia CR, Lammerding J: **Squish and squeeze-the nucleus as a physical barrier during migration in confined environments.** *Curr Opin Cell Biol* 2016, **40**:32-40.
  26. Dittmer TA, Misteli T: **The lamin protein family.** *Genome Biol* 2011, **12**:222.
  27. Torvaldson E, Kochin V, Eriksson JE: **Phosphorylation of lamins determine their structural properties and signaling functions.** *Nucleus* 2015, **6**:166-171.
  28. Dechat T, Gesson K, Foisner R: **Lamina-independent lamins in the nuclear interior serve important functions.** *Cold Spring Harb Symp Quant Biol* 2010, **75**:533-543.
  29. Zwerger M, Roschitzki-Voser H, Zbinden R, Denais C, Herrmann H, Lammerding J, Grutter MG, Medalia O: **Altering lamina assembly reveals lamina-dependent and -independent functions for A-type lamins.** *J Cell Sci* 2015, **128**:3607-3620.
  30. Maurer M, Lammerding J: **The driving force: nuclear mechanotransduction in cellular function, fate, and disease.** *Annu Rev Biomed Eng* 2019, **21**:443-468.
  31. Athirasala A, Hirsch N, Buxboim A: **Nuclear mechanotransduction: sensing the force from within.** *Curr Opin Cell Biol* 2017, **46**:119-127.
  32. Buxboim A, Swift J, Irianto J, Spinler KR, Dingal PC, Athirasala A, Kao YR, Cho S, Harada T, Shin JW *et al.*: **Matrix elasticity regulates lamin-A,C phosphorylation and turnover with feedback to actomyosin.** *Curr Biol* 2014, **24**:1909-1917.
  33. Klapper M, Exner K, Kempf A, Gehrig C, Stuurman N, Fisher PA, Krohne G: **Assembly of A- and B-type lamins studied in vivo with the baculovirus system.** *J Cell Sci* 1997, **110**:2519-2532.
  34. Stuurman N, Heins S, Aebi U: **Nuclear lamins: their structure, assembly, and interactions.** *J Struct Biol* 1998, **122**:42-66.
  35. Makarov AA, Zou J, Houston DR, Spanos C, Solovyova AS, Cardenal-Peralta C, Rappsilber J, Schirmer EC: **Lamin A molecular compression and sliding as mechanisms behind nucleoskeleton elasticity.** *Nat Commun* 2019, **10**:3056.
- Using chemical crosslinking in conjunction with mass-spectrometry, the organization of lamin dimers are dependent on the chemical environment.
36. Zwerger M, Medalia O: **From lamins to lamina: a structural perspective.** *Histochem Cell Biol* 2013, **140**:3-12.
  37. Herrmann H, Aebi U: **Intermediate filaments: molecular structure, assembly mechanism, and integration into functionally distinct intracellular Scaffolds.** *Annu Rev Biochem* 2004, **73**:749-789.
  38. Kapinos LE, Burkhard P, Herrmann H, Aebi U, Strelkov SV: **Simultaneous formation of right- and left-handed anti-parallel coiled-coil interfaces by a coil2 fragment of human lamin A.** *J Mol Biol* 2011, **408**:135-146.
  39. Dhe-Paganon S, Werner ED, Chi YI, Shoelson SE: **Structure of the globular tail of nuclear lamin.** *J Biol Chem* 2002, **277**:17381-17384.
  40. Krimm I, Ostlund C, Gilquin B, Couprie J, Hossenlopp P, Mornon JP, Bonne G, Courvalin JC, Worman HJ, Zinn-Justin S: **The Ig-like structure of the C-terminal domain of lamin A/C, mutated in muscular dystrophies, cardiomyopathy, and partial lipodystrophy.** *Structure* 2002, **10**:811-823.
  41. Strelkov SV, Schumacher J, Burkhard P, Aebi U, Herrmann H: **Crystal structure of the human lamin A coil 2B dimer: implications for the head-to-tail association of nuclear lamins.** *J Mol Biol* 2004, **343**:1067-1080.



42. Ahn J, Jo I, Kang SM, Hong S, Kim S, Jeong S, Kim YH, Park BJ, Ha NC: **Structural basis for lamin assembly at the molecular level.** *Nat Commun* 2019, **10**:3757.  
Structural determination of the lamin A fragment (amino acids 1-285) indicated that the linker domain exhibits a helical conformation.
43. Lilina AV, Chernyatina AA, Guzenko D, Strelkov SV: **Lateral A11 type tetramerization in lamins.** *J Struct Biol* 2019; 107404.  
Structural determination of the helix 1 domain of lamin A, indicated the possible interactions between lateral associated dimers.
44. Foeger N, Wiesel N, Lotsch D, Mucke N, Kreplak L, Aebi U, Gruenbaum Y, Herrmann H: **Solubility properties and specific assembly pathways of the B-type lamin from *Caenorhabditis elegans*.** *J Struct Biol* 2006, **155**:340-350.
45. Wiesel N, Mattout A, Melcer S, Melamed-Book N, Herrmann H, Medalia O, Aebi U, Gruenbaum Y: **Laminopathic mutations interfere with the assembly, localization, and dynamics of nuclear lamins.** *Proc Natl Acad Sci U S A* 2008, **105**:180-185.
46. Ben-Harush K, Wiesel N, Frenkiel-Krispin D, Moeller D, Soreq E, Aebi U, Herrmann H, Gruenbaum Y, Medalia O: **The supramolecular organization of the *C. elegans* nuclear lamin filament.** *J Mol Biol* 2009, **386**:1392-1402.
47. Lucic V, Forster F, Baumeister W: **Structural studies by electron tomography: from cells to molecules.** *Annu Rev Biochem* 2005, **74**:833-865.
48. Irobalieva RN, Martins B, Medalia O: **Cellular structural biology as revealed by cryo-electron tomography.** *J Cell Sci* 2016, **129**:469-476.
49. Weber MS, Wojtynek M, Medalia O: **Cellular and structural studies of eukaryotic cells by Cryo-electron tomography.** *Cells* 2019, **8**.
50. Elad N, Volberg T, Patla I, Hirschfeld-Warneken V, Grashoff C, Spatz JP, Fassler R, Geiger B, Medalia O: **The role of integrin-linked kinase in the molecular architecture of focal adhesions.** *J Cell Sci* 2013, **126**:4099-4107.
51. Sorrentino S, Studt JD, Horev MB, Medalia O, Sapra KT: **Toward correlating structure and mechanics of platelets.** *Cell Adhes Migr* 2016, **10**:568-575.
52. Medalia O, Weber I, Frangakis AS, Nicastro D, Gerisch G, Baumeister W: **Macromolecular architecture in eukaryotic cells visualized by cryoelectron tomography.** *Science* 2002, **298**:1209-1213.
53. Harapin J, Eibauer M, Medalia O: **Structural analysis of supramolecular assemblies by cryo-electron tomography.** *Structure* 2013, **21**:1522-1530.
54. Medeiros JM, Bock D, Pilhofer M: **Imaging bacteria inside their host by cryo-focused ion beam milling and electron cryotomography.** *Curr Opin Microbiol* 2018, **43**:62-68.
55. Ghosal D, Kim KW, Zheng H, Kaplan M, Truchan HK, Lopez AE, McIntire IE, Vogel JP, Cianciotto NP, Jensen GJ: **In vivo structure of the *Legionella* type II secretion system by electron cryotomography.** *Nat Microbiol* 2019, **4**:2101-2108.
56. Baker LA, Grange M, Grunewald K: **Electron cryo-tomography captures macromolecular complexes in native environments.** *Curr Opin Struct Biol* 2017, **46**:149-156.
57. Albert S, Wietrzynski W, Lee CW, Schaffer M, Beck F, Schuller JM, Salome PA, Plitzko JM, Baumeister W, Engel BD: **Direct visualization of degradation microcompartments at the ER membrane.** *Proc Natl Acad Sci U S A* 2019.
58. Turgay Y, Eibauer M, Goldman AE, Shimi T, Khayat M, Ben-Harush K, Dubrovsky-Gaupp A, Sapra KT, Goldman RD, Medalia O: **The molecular architecture of lamins in somatic cells.** *Nature* 2017, **543**:261-264.  
At the lamina, lamin filaments are 300-400 nm long and ~3.5 nm in crosssection, as revealed by cryo-electron tomography.
59. Zingerman-Koladko I, Khayat M, Harapin J, Shoseyov O, Gruenbaum Y, Salman A, Medalia O, Ben-Harush K: **The assembly of *C. elegans* lamins into macroscopic fibers.** *J Mech Behav Biomed Mater* 2016, **63**:35-43.
60. Sapra T, Qin Z, Buehler MJ, Aebi U, Dubrovsky-Gaupp A, Mueller DJ, Medalia O: **Nonlinear mechanics of lamin filaments and the meshwork topology build an emergent nuclear lamina.** *bioRxiv* 2019:846550.  
A direct mechanical measurements of lamin LIII filaments.
61. Shimi T, Kittisopikul M, Tran J, Goldman AE, Adam SA, Zheng Y, Jaqaman K, Goldman RD: **Structural organization of nuclear lamins A, C, B1, and B2 revealed by superresolution microscopy.** *Mol Biol Cell* 2015, **26**:4075-4086.
62. Nmezi B, Xu J, Fu R, Armiger TJ, Rodriguez-Bey G, Powell JS, Ma H, Sullivan M, Tu Y, Chen NY *et al.*: **Concentric organization of A- and B-type lamins predicts their distinct roles in the spatial organization and stability of the nuclear lamina.** *Proc Natl Acad Sci U S A* 2019, **116**:4307-4315.  
The position of the highest concentration of lamin A is 5 nm into the nucleoplasm in comparison the that of lamin B, as revealed by super resolution microscopy.
63. Shevelyov YY, Ulianov SV: **The nuclear lamina as an organizer of chromosome architecture.** *Cells* 2019, **8**.
64. Romero-Bueno R, de la Cruz Ruiz P, Artal-Sanz M, Askjaer P, Dobrzynska A: **Nuclear organization in stress and aging.** *Cells* 2019, **8**.
65. Guerreiro I, Kind J: **Spatial chromatin organization and gene regulation at the nuclear lamina.** *Curr Opin Genet Dev* 2019, **55**:19-25.
66. Kind J, Pagie L, de Vries SS, Nahidiazar L, Dey SS, Bienko M, Zhan Y, Lajoie B, de Graaf CA, Amendola M *et al.*: **Genome-wide maps of nuclear lamina interactions in single human cells.** *Cell* 2015, **163**:134-147.
67. Brunet A, Forsberg F, Fan Q, Saether T, Collas P: **Nuclear Lamin B1 Interactions with chromatin during the circadian cycle are uncoupled from periodic gene expression.** *Front Genet* 2019, **10**:917.
68. Marko M, Hsieh C, Schalek R, Frank J, Mannella C: **Focused-ion-beam thinning of frozen-hydrated biological specimens for cryo-electron microscopy.** *Nat Methods* 2007, **4**:215-217.
69. Wagenknecht T, Hsieh C, Marko M: **Skeletal muscle triad junction ultrastructure by focused-ion-beam milling of muscle and cryo-electron tomography.** *Eur J Transl Myol* 2015, **25**:4823.
70. Gorelick S, Buckley G, Gervinskis G, Johnson TK, Handley A, Caggiano MP, Whisstock JC, Pocock R, de Marco A: **PIE-scope, integrated cryo-correlative light and FIB/SEM microscopy.** *eLife* 2019, **8**.
71. Schaffer M, Pfeffer S, Mahamid J, Kleindiek S, Laugs T, Albert S, Engel BD, Rummel A, Smith AJ, Baumeister W *et al.*: **A cryo-FIB lift-out technique enables molecular-resolution cryo-ET within native *Caenorhabditis elegans* tissue.** *Nat Methods* 2019, **16**:757-762.  
The lift-out approach opens large number of possibilities to acquire structural information out off vitrified tissues.
72. Harapin J, Bormel M, Sapra KT, Brunner D, Kaech A, Medalia O: **Structural analysis of multicellular organisms with cryo-electron tomography.** *Nat Methods* 2015, **12**:634-636.
73. Cai S, Bock D, Pilhofer M, Gan L: **The in situ structures of mono-, di-, and trinucleosomes in human heterochromatin.** *Mol Biol Cell* 2018, **29**:2450-2457.
74. Mahamid J, Pfeffer S, Schaffer M, Villa E, Danev R, Cuellar LK, Forster F, Hyman AA, Plitzko JM, Baumeister W: **Visualizing the molecular sociology at the HeLa cell nuclear periphery.** *Science* 2016, **351**:969-972.
75. Mosalaganti S, Kosinski J, Albert S, Schaffer M, Strenkert D, Salome PA, Merchant SS, Plitzko JM, Baumeister W, Engel BD *et al.*: **In situ architecture of the algal nuclear pore complex.** *Nat Commun* 2018, **9**:2361.



76. Pfeifer CR, Irianto J, Discher DE: **Nuclear mechanics and cancer cell migration.** *Adv Exp Med Biol* 2019, **1146**:117-130.
  77. Zwerger M, Ho CY, Lammerding J: **Nuclear mechanics in disease.** *Annu Rev Biomed Eng* 2011, **13**:397-428.
  78. Stephens AD, Banigan EJ, Adam SA, Goldman RD, Marko JF: **Chromatin and lamin A determine two different mechanical response regimes of the cell nucleus.** *Mol Biol Cell* 2017, **28**:1984-1996.
  79. Wintner O, Hirsch-Attas N, Schlossberg M, Brofman F, Friedman R, Kupervaser M, Kitsberg D, Buxboim A: **A unified linear viscoelastic model of the cell nucleus defines the mechanical contributions of lamins and chromatin.** *Adv Sci (Weinh)* 2020, **7**:1901222.
- While lamin-A and lamin-B1 contribute to nucleus stiffness, viscosity is specified mostly by lamin-A. Elastic and viscous association of lamin-B1 and lamin-A is supported by transcriptional and proteomic profiling analyses.
80. Aebersold U, Cohn J, Buhle L, Gerace L: **The nuclear lamina is a meshwork of intermediate-type filaments.** *Nature* 1986, **323**:560-564.